

FOR ANIMAL USE ONLY

Pegaforte®

Reg. No. G3768 Act 36/1947

[NSO] V07/18.1.8/447 Act 13/2003

ORAL PASTE

Apple flavour

CAUTION

INDICATIONS

Broad spectrum roundworm, tapeworm and bot remedy for horses of all ages including pregnant mares.

COMPOSITION

Ivermectin 12 mg/g

Praziquantel 150 mg/g

STORAGE

Store in a cool, dry place below 25 °C.

WARNINGS

- DO NOT USE ON ANIMALS INTENDED FOR HUMAN CONSUMPTION.
- Pregnant mares should be treated with the usual additional care.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Store in a safe place.
- Do not smoke or eat while handling the product.
- Wash hands after use.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- **Pegaforte®** is administered orally.
- Each marking on the syringe delivers enough paste for 100 kg body mass.
- The recommended dosage is 200 µg of Ivermectin and 2,5 mg of Praziquantel per kg of body mass.
- One 10 g syringe contains sufficient paste to treat one 600 kg horse at the recommended dose rate.

ADMINISTRATION

- The body mass of the horse should be determined accurately to facilitate administration of the correct dose.
- Before administration, ensure that the horse does not have food in its mouth.
- Insert syringe in the gap between the front and back teeth and deliver paste onto the back of the tongue.
- Immediately raise the horse's head for a few seconds after dosing to ensure swallowing of the paste.

EFFICACY

Pegaforte® broad spectrum parasiticide effectively controls the following worms in horses:

Large strongyles <i>Strongylus vulgaris</i> (Bloodworm) <i>Strongylus edentatus</i> <i>Strongylus equinus</i> <i>Triodontophorus</i>	Adults & arterial larvae Adults & encysted larvae tissue stage Adults Adults	Hairworms <i>Trichostrongylus axei</i>	Adults
Small strongyles <i>Cyathostomum</i> spp. <i>Cylicocyclus</i> spp. <i>Cylicostephanus</i> spp. <i>Cylicodontophorus</i> spp. <i>Gyalocephalus</i> spp.	Adults & 4th stage larvae	Intestinal threadworms <i>Strongyloides westeri</i>	Adults
Ascarids <i>Parascaris equorum</i>	Adults and 3rd & 4th stage larvae	Large-mouthed stomach worms <i>Habronema muscae</i> <i>Draschia</i> spp.	Adults 3rd stage larvae
Pinworms <i>Oxyuris equi</i>	Adults and 3rd & 4th stage larvae	Lungworms <i>Dictyocaulus arnfieldi</i>	Adults & 4th stage larvae
		Bots <i>Gasterophilus</i> spp.	Oral and gastric stages
		Cestodes <i>Anoplocephala</i> spp.	

Pegaforte® is also effective for the control of skin lesions caused by *Habronema* spp. and *Draschia* spp. cutaneous larvae (summer sores) and *Onchocerca* spp. microfilariae (cutaneous onchocerciasis).

PROPOSED WORM CONTROL PROGRAMME

Horses of all ages, including pregnant mares, breeding stallions and foals can be treated with **Pegaforte®**.

Mares: Regular dosing at 6-8 week intervals on pastures during summer and autumn.

Stallions: Treat at least twice during the year.

Foals: Initial treatment at ± 2 months of age and followed by strategic treatment at 6-8 week intervals.

Bots: Bot flies are active during late spring and peak during autumn. The bots are then carried through the winter. Treatment against the bots that are attached to the stomach wall is essential in order to remove the established burdens in the autumn and winter.

Consult your veterinarian for a worm control programme.



Registration Holder:
Kyron Vet Rx (Pty) Ltd
 Co. Reg. No. 2001/017471/07
 41 Park Avenue
 Highway Business Park
 Rooihuiskraal Ext 31
 Centurion, 0157, South Africa

SLEGS VIR DIEREGEBRUIK

Pegaforte®

Reg. Nr. G3768 Wet 36/1947

[NSO] V07/18.1.8/447 Act 13/2003

ORALE PASTA

Appel geur

VERSIGTIG

AANWYSINGS

Breë-spektrum middel teen rondewurms, lintwurms en papies in perde van alle ouderdomme, insluitend dragtige merries.

SAMESTELLING

Ivermektien 12 mg/g

Prasikwantel 150 mg/g

BERGING

Berg in 'n koel, droë plek benede 25 °C.

WAARSKUWINGS

- MOENIE DIERE BEHANDEL WAT VIR MENSLIKE VERBRUIK BESTEM IS NIE.
- Dragtige merries moet soos gewoonlik met addisionele sorg behandel word.
- Hou buite bereik van oningeligte persone, kinders en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMATREËLS

- Berg in 'n veilige plek.
- Moenie rook of eet tydens hantering van die produk nie.
- Was hande na gebruik.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- **Pegaforte®** word oraal toegedien.
- Elke merk op die spuit dien genoeg pasta toe vir 100 kg liggaamsmassa.
- Die aanbevole dosis is 200 µg Ivermektien en 2,5 mg Prasikwantel per kg liggaamsmassa.
- Een 10 g spuit bevat genoeg pasta om een 600 kg perd teen die aanbevole dosis te behandel.

TOEDIENING

- Die perd se liggaamsmassa moet akkuraat bepaal word om die korrekte dosis te verseker.
- Voor toediening, maak seker dat daar geen kos in die perd se bek is nie.
- Plaas die spuit in die kant van die perd se bek in die gaping tussen die voor- en agtertande en deponeer die pasta agter op die tong.
- Lig onmiddellik die perd se kop vir 'n paar sekondes om te verseker dat die volle dosis ingesluk word.

DOELTREFFENDHEID

Pegaforte® breë-spektrum parasietmiddel bied effektiewe beheer van die volgende wurms in perde:

Grootstrongiele <i>Strongylus vulgaris</i> (Bloedwurm) <i>Strongylus edentatus</i>	Volwassenes & slagaar larwes	Haarwurms <i>Trichostrongylus axei</i>	Volwassenes
<i>Strongylus equines</i> <i>Triodontophorus</i>	Volwassenes & ingesulpte larwes – weefsel stadium	Ingewands draadwurms <i>Strongyloides westeri</i>	Volwassenes
Kleinstrongiele <i>Cyathostomum</i> spp. <i>Cylicocyclus</i> spp. <i>Cylicostephanus</i> spp. <i>Cylicodontophorus</i> spp. <i>Gyalocephalus</i> spp.	Volwassenes & 4de stadium larwes	Grootbekmaagwurms <i>Habronema muscae</i> <i>Draschia</i> spp.	Volwassenes 3de stadium larwes
Askariedes <i>Parascaris equorum</i>	Volwassenes en 3de & 4de stadium larwes	Longwurms <i>Dictyocaulus arnfieldi</i>	Volwassenes & 4de stadium larwes
Sambokwurms <i>Oxyuris equi</i>	Volwassenes en 3de & 4de stadium larwes	Papies <i>Gasterophilus</i> spp.	Orale en maag stadia
		Sestodes <i>Anoplocephala</i> spp.	

Pegaforte® is ook doeltreffend vir die beheer van velletsels veroorsaak deur *Habronema* spp. en *Draschia* spp. vellarwes (somer sere) en *Onchocerca* spp. mikrofilariae (liggamentwurm).

VOORGESTELDE WURMBEHEER PROGRAM

Perde van alle ouderdomme, insluitend dragtige merries, stoethingste en vullens kan met **Pegaforte®** behandel word.

Merries: Gereelde dosering met 6-8 week tussenposes gedurende somer en herfs.

Stoethingste: Behandel ten minste 2 keer per jaar.

Vullens: Eerste behandeling op ± 2 maande ouderdom gevolg deur strategiese behandeling met 6-8 week tussenposes.

Papies: Papievlieë is aktief gedurende laat lente en piek tydens herfs. Die papies word deur die winter gedra. Behandeling teen papies aan die maagwand is noodsaaklik om gevestigde laste in die herfs en winter te verwyder.

Raadpleeg u veearts vir 'n wurmbesluiting program.